




XX ITALIAN-SPANISH CONGRESS ON THERMODYNAMICS OF METAL COMPLEXES

XXXVI ANNUAL CONGRESS OF THE “GRUPPO DI TERMODINAMICA DEI COMPLESSI”



Tirrenia (Pisa), 7 -11 June 2009

SCIENTIFIC COMMITTEE

Antonio Bianchi, Università di Firenze
Guido Crisponi, Università di Cagliari
Enrique García España, Universitat de Valencia
Juan Niclós Gutiérrez, Universidad de Granada
José Maria Leal, Universidad de Burgos
Francesco Malatesta, Università di Pisa
Maria Angeles Olazabal, Universidad del País Vasco
Silvio Sammartano, Università di Messina
Francisco Sánchez Burgos, Universidad de Sevilla
Fernando Secco, Università di Pisa
Maria Rosaria Tiné, Università di Pisa
Manuel Valiente, Universitat Autònoma de Barcelona
Marcella Venturini, Università di Pisa

ORGANIZING COMMITTEE

Fernando Secco (Chairman)
Tarita Biver (Secretary)
Marcella Venturini
Alessia Boggioni
Sabriye Aydinoglu
Maria Rosa Beccia
Nurettin Eltugral
Lisa Ghezzi
Elisa Turriani

SECRETARIAT

Tarita Biver
Univeristà di Pisa
Dipartimento di Chimica e Chimica Industriale
Via Risorgimento 35 - 56126 Pisa (Italy)

Tel./ Fax. +39 050 2219259/260
e-mail: tarita@dccci.unipi.it

THE INTERACTION OF NATIVE DNA WITH Ni(II), Cu(II) and Zn(II) COMPLEXES OF 5-TRIETHYL AMMONIUM METHYL SALICYLIDENE ORTHO-PHENYLENDIIMINE

Giampaolo Barone^a, Angela Ruggirello^b, Arturo Silvestri^a, Alessio Terenzi^a, Vincenzo Turco Liveri^b

^a*Dipartimento di Chimica Inorganica e Analitica "S. Cannizzaro" and ^bDipartimento di Chimica Fisica "F. Accascina", Università di Palermo, Viale delle Scienze, Parco d'Orleans II, Edificio 17, 90128 Palermo (Italy), e-mail: gbarone@unipa.it*

This communication will give an account of recent studies on the interaction of native calf thymus DNA with the cationic Ni(II), Cu(II) and Zn(II) complexes of 5-triethyl ammonium methyl salicylidene ortho-phenylendiimine (H_2L^{2+}), see Figure 1, in 1 mM Tris-HCl aqueous solutions at neutral pH. The title interaction has been monitored as a function of the metal complex-DNA molar ratio by UV absorption spectrophotometry, circular dichroism (CD) and fluorescence spectroscopy [1,2].




Figure 1: Structure of the ML^{2+} complexes ($M = Ni, Cu, Zn$)

The remarkable modifications of the DNA CD spectrum, the appearance of a broad induced CD band in the range 350-450 nm, the strong increase of the DNA melting temperature (T_m), the fluorescence quenching of ethidium bromide-DNA solutions in the presence of increasing amounts of the metal complexes, and the high values of the intrinsic binding constants (K_b) determined by UV spectrophotometric titrations, support the existence of a tight intercalative interaction of the three metal complexes with DNA.

The interaction of CuL^{2+} with native DNA has been also investigated in reverse micelles, constituting nanoscopic domains biomimetic of the intra-cellular environment [3]. The results of this study indicate the occurrence of dramatic structural changes of both the DNA and the metal complex-DNA system, when going from aqueous solution to the micellar phase [3].

The interaction of ZnL^{2+} with DNA has been recently revisited, in consideration of the photophysical processes induced on the zinc complex by visible light. Steady state and time-resolved fluorescence spectroscopy measurements, supported by quantum chemical calculations, point out that ZnL^{2+} is effectively protected by DNA from photoinduced redox phenomena, which occur when it is in the free state dispersed in aqueous buffer solution.

[1] G. Barone et al., J. Inorg. Biochem., in press, doi:10.1016/j.jinorgbio.2009.01.006.

[2] A. Silvestri et al., J. Inorg. Biochem., **2007**, 101, 841-848.

[3] G. Barone et al., Dalton Trans., **2008**, 4172-4178.